

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human NKp80/KLRF1 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human (rh) NKp30, rhNKp44, rhNKp46, or recombinant mouse NKp46 is observed.
Source	Monoclonal Rat IgG _{2A} Clone # 239127
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human NKp80/KLRF1 Val66-Tyr231 Accession # Q9NZS2
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	Human whole blood CD56 ⁺ natural killer cells

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

NKp80, also known as killer cell lectin-like receptor subfamily F member 1 (KLRF1), is an 80 kDa type II transmembrane, homodimeric, activating co-receptor expressed exclusively on the surface of NK cells. Human NKp80, a member of the C-type lectin family of proteins (1-3), is 231 aa in length with a 38 amino acid (aa) cytoplasmic region, a 21 aa transmembrane region, and a 172 aa extracellular domain (ECD). The ECD contains the C-type lectin-like domain (aa 114-230) and four potential sites of N-linked glycosylation. Multiple splicing variants produce four isoforms for NKp80. Isoform 1 is the standard protein. There are no murine orthologs for NKp80 (4). NKp80 functions as an inhibitory receptor based on the presence of two immunoreceptor tyrosine-based inhibitory motifs (ITIM) in the N-terminal cytoplasmic region and the lack of charged amino acids in the transmembrane region (1). However, later studies show that NKp80 displays an activating function rather than an inhibitory one (2-3, 5). In *in vitro* experiments in human NK cells, NKp80 stimulates NK cell cytotoxicity and induces calcium influx after triggering by appropriate antibodies (2, 3). One study reports that NKp80 likely functions as a co-receptor, cooperating with NKp46 and other activating receptors in NK cell cytotoxicity (2). Activation-induced C-type lectin (AICL) is the ligand for NKp80 (3). It is a myeloid-specific receptor expressed by monocytes, macrophages, and granulocytes (3). Crosslinking of both NKp80 and AICL stimulates secretion of proinflammatory cytokines suggesting that NKp80-AICL interaction is involved in the activating crosstalk between NK cells and myeloid cells, and thus may influence the initiation and maintenance of immune responses in humans (3).

References:

1. Roda-Navarro, P. *et al.* (2000) *Eur. J. Immunol.* **30**:568.
2. Vitale, M. *et al.* (2001) *Eur. J. Immunol.* **31**:233.
3. Welte, S. *et al.* (2006) *Nat. Immunol.* **7**:1334.
4. Yokoyama, W.M. and B.F. Plougastel (2003) *Nat. Rev. Immunol.* **3**:304.
5. Thomas, M. *et al.* (2008) *Proc. Natl. Acad. Sci. U.S.A.* **105**:1656.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc. and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.