

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human SP-D in direct ELISAs.
Source	Monoclonal Mouse IgG _{2B} Clone # 292215
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human SP-D Ala21-Phe375 Accession # P35247.2
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

ELISA Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

SP-D (surfactant protein-D; also PSP-D) is a 43 kDa member of the collectin family of innate immune modulators. It is constitutively secreted by alveolar lining cells and epithelium associated with tubular structures. Its principal components consist of a collagen-like region and a C-terminal carbohydrate recognition domain (CRD), a structure that further places it in a subset of an expanded group of proteins termed defense collagens (1-4). Human SP-D is synthesized as a 375 amino acid (aa) precursor. It contains a 20 aa signal sequence and a 355 aa mature region. The mature molecule is characterized by the presence of a 25 aa N-terminal linking-region, a 177 aa hydroxyproline and hydroxylysine collagen-like domain, a 46 aa coiled-coil segment, and a 106 aa, C-terminal collectin-like C-type lectin domain (CRD) (5, 6). Two additional, potential isoforms exist. One shows a 13 aa N-terminal extension, while the other combines the N-terminal extension with a deletion of aa's 206-375. Mature human SP-D shares 75% and 78% aa identity with mouse and pig SP-D, respectively. Monomeric SP-D is unusual (3). The basic form of SP-D is that of a glycosylated, disulfide-linked 150 kDa trimer that generates an α -helical coiled-coil structure linked to a "head" of three symmetrical CRDs (4, 7). Each CRD recognizes the hydroxides of one monosaccharide (4). Trimerization allows for the discrimination of monosaccharide patterns specific to microbial pathogens (7). Typically, SP-D forms a higher-order 620 kDa, X-shaped dodecamer through disulfide bonds associated with the N-terminus (8). This allows for even finer discrimination of self vs. nonself carbohydrate patterns, and facilitates binding to complex antigens (8, 9). One polymorphism, a Met11-Thr11 transition in human, apparently precludes the formation of oligomers, potentially affecting the ability of affected individuals to interact with microorganisms (9, 10). Finally, SP-D is known to bind both SIRP α and the calreticulin/CD91 complex on macrophages. When the ratio of antigen/pathogen to available CRDs is low, antigen can be bound without occupying all available CRDs. The free CRDs will bind to SIRP α , generating a signal that downmodulates the inflammatory response. When virtually all CRDs are occupied by ligand, however, free CRDs are not available for SIRP α binding. Instead, the dodecamer is depicted to undergo a structural rearrangement, exposing the N-termini of all four linked trimers. This exposed terminus is known to bind to the calreticulin/CD91 complex, an event that initiates inflammation. Thus, it would appear that SP-D allows for a graded response to environmental challenge. SP-D provides a mechanism for the clearance of small antigenic insults without the need for a damaging inflammatory response (3).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.