

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human BMP-1/PCP in direct ELISA.
Source	Monoclonal Rat IgG _{2A} Clone # 264801
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human BMP-1/PCP Ala121-Gln730 Accession # NP_001190
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunohistochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Bone morphogenetic protein 1 (BMP-1), also known as procollagen C-proteinase (PCP), is a zinc protease of the astacin family (1, 2). BMP-1/PCP plays a key role in formation of extracellular matrix (ECM) by converting precursor proteins into their mature and functional forms. The precursor proteins identified as substrates for BMP-1/PCP include collagens, biglycan, laminin 5, dentin matrix protein-1, and lysyl oxidase (3). There are six alternatively spliced forms known to be derived from the BMP-1 gene, and isoform 1 consisting of residues 1 to 730 was expressed. The secreted and purified protein does not contain the signal peptide (amino acid residues 1-22) and pro domain (residues 23-120), but contain protease (residues 121-321), CUB I (residues 322-434), CUB II (residues 435-546), EGF-like (residues 547-588) and CUB III (residues 591-703) domains. The pro domain is apparently cleaved by a furin-like proprotein convertase (4). The purified BMP-1/PCP is an active protease and its peptidase activity can be determined as described above. The purified BMP-1/PCP is predicted to possess procollagen C-proteinase activity because it contains the minimal domain structure required (5).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.