

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human ICAM-5 in direct ELISAs and Western blots. In Western blots, no cross-reactivity with recombinant human (rh) CD31, rhCEACAM-1, recombinant mouse (rm) DCC, rhICAM-1, -2, -3, rmlCAM-5, rhMAdCAM-1, or rmVCAM-1 is observed.
Source	Monoclonal Mouse IgG _{2A} Clone # 283607
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human ICAM-5 Ala28-Glu570 Accession # Q9UMF0
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

ELISA Capture (Matched Antibody Pair)	Optimal dilution of this antibody should be experimentally determined.
ELISA Detection (Matched Antibody Pair)	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Intercellular adhesion molecule-5 (ICAM-5), also known as telencephalin, is a cell surface glycoprotein belonging to the immunoglobulin superfamily. Human ICAM-5 consists of an 832 amino acid (aa) extracellular domain containing 9 immunoglobulin (Ig) domains and 15 N-glycosylation sites, a 28 aa transmembrane domain, and a 64 aa cytoplasmic domain. ICAM-5 shares 38-55% aa identity with other ICAMs, being most closely related to ICAM-1 (50% identity) and ICAM-3 (55% identity) (1). Human and mouse ICAM-5 share 85% aa identity. The tissue distribution of ICAM-5 is unique among ICAMs, being expressed only in telencephalic regions of the central nervous system (2). Like other ICAMs, ICAM-5 binds to the leukocyte integrin LFA-1 (CD11a/CD18) (3). Binding of ICAM-5 to LFA-1 is dependent on the first amino terminal Ig domain of ICAM-5 (4). ICAM-5 also displays homophilic binding, with the amino terminal Ig domain binding to Ig domains 4-5. Homophilic binding of ICAM-5 is dependent of ICAM-5 being in a monomeric form. The monomeric form of ICAM-5 is found during dendritogenesis in developing brain, whereas a high molecular weight complex is found in mature neurons (5).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.