

DESCRIPTION

Species Reactivity	Canine
Specificity	Detects canine KGF/FGF-7 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human (rh) FGF-11, -12, -13, -16, -20, -21, -22, -23, or rhFGFbasic is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 299026
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant canine KGF/FGF-7 Cys32-Thr194 Accession # P79150
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

The fibroblast growth factor (FGF) family consists of twenty three known structurally-related proteins that play key roles in development, morphogenesis, angiogenesis, wound healing and tumorigenesis (1). All FGFs share a common structural fold, the β-trefoil scaffold, in a conserved central region of approximately 120 amino acid (aa), which contains the active site and a heparin binding site (2). FGF-7, also known as keratinocyte growth factor (KGF), was originally identified from the conditioned medium of a human embryonic lung fibroblast cell line (3). It is secreted by cells of mesenchymal origin and functions predominantly on epithelial cells (1). The mature caFGF-7 is generated from a 194 aa precursor protein by cleavage of the first 31 aa at the N-terminus. It shares 97.4% and 95.9% sequence identity with human and mouse KGF, respectively (4). FGFs signal through high affinity tyrosine kinase receptors (FGF R) (5). Cellular responses to FGFs are modulated by heparan sulfate proteoglycans that are also known as low affinity FGF receptors and by heparin (6). Various isoforms of the four FGF R exist (FGF R-1 through -4). They are produced from alternatively spliced transcripts in both the extracellular and intracellular domains. FGF-7 signals through the FGF R2IIIb (also known as KGF R), which is a spliced variant of FGF R2 (7). KGF R binds FGF-1 and FGF-7 with high affinity and FGF-2 with a 20-fold lower affinity (8).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.