

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human IL-24 in ELISAs.
Source	Monoclonal Mouse IgG _{2A} Clone # 283161
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human IL-24 Gln50-Leu206 Accession # Q13007
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

ELISA Capture (Matched Antibody Pair)	Optimal dilution of this antibody should be experimentally determined.
ELISA Detection (Matched Antibody Pair)	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Interleukin 24 (IL-24), also known as mda-7 (melanoma differentiation associated gene-7), is a member of the IL-10 family of helical cytokines. The IL-24 gene encodes a precursor protein of 207 amino acids (aa) that contains a 48 aa signal sequence and an 18 kDa, 158 aa mature segment. There are three potential N-linked glycosylation sites, at least one of which is used. When secreted, IL-24 is a 35-40 kDa phosphorylated glycoprotein that apparently can exist as either a monomer or dimer. It is suggested that glycosylation is essential for activity. Mature human IL-24 shares 69% aa sequence identity with mouse and rat IL-24. Human IL-24 is also active in rodent systems. Cells known to express IL-24 include B cells, CD4⁺ T cells, NK cells, lymph node dendritic cells, monocytes, melanocytes, and melanoma cells. Functionally, IL-24 has diverse activities. At low concentrations on monocytes, it induces type I proinflammatory cytokines such as IFN-γ, IL-1β, IL-12, and TNF-α. At high concentrations, it is a strong inducer of apoptosis in tumor cells, but not normal cells. IL-24 also has anti-angiogenic properties. It directly binds IL-24 receptors on endothelial cells, activating STAT3 and blocking their differentiation. IL-24 binds and signals through two heterodimeric receptor complexes. One complex is the combination of IL-20 Rα and IL-20 Rβ, which is shared with IL-19 and IL-20. The second complex is a combination of IL-22 R and IL-20 Rβ, which is shared with IL-20.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.