

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human LILRB1/CD85j/ILT2 in direct ELISAs and Western blots. In direct ELISAs, no cross-reactivity with recombinant human (rh) ILT1, 3, 4, 5, rhLIR-6 or rhLIR-8 is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 292319
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human LILRB1/CD85j/ILT2 Gly24-His458 Accession # Q8NHL6
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

CyTOF-reported	Optimal dilution of this antibody should be experimentally determined.
Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

The immunoglobulin-like transcript (ILT) family of activating and inhibitory type immunoreceptors are expressed on many leukocyte subsets and function in the regulation of immune responses (1-3). This family was also named leukocyte Ig-like receptors (LIR) and monocyte/macrophage Ig-like receptors (MIR). ILTs share significant homology with killer cell Ig-like receptors (KIR). The ILT genes are located on human chromosome 19q13.4 in the leukocyte receptor complex, which also include the genes encoding KIRs (4). With the exception of ILT-6, which is a soluble molecule, all ILT family members are type I transmembrane proteins having two or four extracellular Ig-like domains (2, 3). One subset of the ILT receptors (referred to as subfamily B of the LIRs) has long cytoplasmic tails containing immunoreceptor tyrosine-based inhibitory motifs (ITIMs) that inhibit signaling events by recruiting SH2-containing protein tyrosine phosphatase-1. Another subset of the ILT receptors (referred to as subfamily A of the LIRs) contains activating receptors with short cytoplasmic regions that lack signal transduction motifs. These receptors contain a basic arginine residue within their transmembrane domains, which allows association with Fc Rγ, an immunoreceptor tyrosine-based activation motif (ITAM)-bearing signal adapter protein (1-3). ILT2, also known as LIR1, MIR7, and CD85j, is expressed on most monocytes, dendritic cells, and mature B cells (1-3). It is also expressed on small percentages of T cells and NK cells. ILT2 has four extracellular Ig-like domains and three cytoplasmic ITIMs. It functions as an inhibitory receptor that prevents cellular activation. ILT2 has been shown to bind classical (HLA-A and -B) and nonclassical (HLA-G1, -E and -F) MHC class I molecules (MHC I) (1-3). ILT2 also binds with high affinity to an MHC class I homologue from human cytomegalovirus (3). Ligation of ILT2 by MHC class I may function to poise cellular activation thresholds and inhibit various leukocyte effector mechanisms that are regulated by MHC class I molecules on target cells.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.