

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Pentraxin 3/TSG-14 in direct ELISAs and Western blots. In direct ELISAs and Western blots, approximately 40-100% cross-reactivity with recombinant human Pentraxin 3 is observed.
Source	Monoclonal Rat IgG _{2B} Clone # 265629
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse Pentraxin 3/TSG-14 Glu18-Ser381 Accession # P48759
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Pentraxin 3 (PTX3), also known as TSG-14, was initially identified as a TNF- α or IL-1 β inducible gene (1-3). It belongs to the pentraxin family, which was named originally for the homo-pentameric structure formed by its members (4). The pentraxin family is divided into two subfamilies: the "short" and the "long" pentraxins with approximate molecular weights of 25 kDa and 50 kDa, respectively. TSG-14 is a member of the long pentraxin subfamily, which also includes the *Xenopus laevis* XL-PXN1, the guinea pig apelin/p50, the rat neuronal pentraxin I (NPI) and NPR, the human neuronal pentraxin II (NPTX2) and the human neuronal activity-related pentraxin (5). Mature secreted PTX3 contains a pentaxin-like domain at its carboxy-terminus that shares 23-28% amino acid (aa) sequence similarity to C-reactive protein (CRP) and serum amyloid P component (SAP), which belong to the short pentraxin subfamily. However, the N-terminal sequence of TSG-14 does not share aa sequence homology with any of the "short" pentaxins (3). Unlike CRP and SAP, which forms pentamers only, TSG-14 forms both pentameric and higher ordered oligomers (5). Similar to CRP and SAP, TSG-14 binds to the complement cascade component C1q (6). However, TSG-14 does not bind to phosphoethanolamine, phosphocholine, or high pyruvate agarose, which are known ligands for CRP and SAP. While CRP and SAP are primarily produced in the liver, TSG-14 expression is strongly upregulated by TNF- α , IL-1 β , and bacterial LPS in peripheral fibroblasts, endothelial cells, and macrophages (7). At the amino acid level, human and mouse TSG-14 share 88% aa sequence homology (8). TSG-14 concentration is elevated in the joint fluid of patients with rheumatoid arthritis (RA), indicating that TSG-14 may be a potential mediator of immune response (9). TSG-14 may also function in the regulation of the uptake and clearance of apoptotic cells by dendritic cells (10). An *in vivo* study showed that TSG-14 transgenic mice are more resistant to sepsis and endotoxemia compared to wild-type during inflammatory injury (11). Increased expression of TSG-14 may enhance the immune response to protect the host from infection.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.