

**DESCRIPTION**


<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse NKp46/NCR1 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human (rh) NKp30, rhNKp44, rhNKp46, or rhNKp80 is observed.
<b>Source</b>	Monoclonal Rat IgG <sub>2A</sub> Clone # 29A1.4
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Murine myeloma cell line NSO-derived recombinant mouse NKp46-Fc chimera Glu22-Asn255 Accession # Q8C567
<b>Conjugate</b>	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
<b>Formulation</b>	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Flow Cytometry</b>	0.25-1 µg/10 <sup>6</sup> cells	See Below

**DATA**



**PREPARATION AND STORAGE**

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Protect from light. Do not freeze.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul>

#### BACKGROUND

NKp46, along with NKp30 and NKp44, are activating receptors that have been collectively termed the natural cytotoxicity receptors (NCR) (1). These receptors are expressed almost exclusively by NK cells and play a major role in triggering some of the key lytic activities of NK cells. In human systems, the CD56<sup>dim</sup>CD16<sup>+</sup> subpopulation that makes up the majority of NK cells in the peripheral blood and spleen expresses NKp46 in both resting and activated states (2). The main NK cell population of the lymph node (CD56<sup>bright</sup>CD16<sup>-</sup>) expresses low levels of NKp46 in resting cells, but expression is upregulated by IL-2. Mouse NKp46, also known as MAR-1 (3), is a type I transmembrane protein with two extracellular Ig-like domains. It has a positive charge in its transmembrane domain that permits association with the ITAM-bearing signal adapter proteins, CD3 $\zeta$  and Fc $\epsilon$  R1 $\gamma$  (4). Studies with neutralizing antibodies indicate that the three NCR are primarily responsible for triggering the NK-mediated lysis of many human tumor cell lines. Blocking any of the NCRs individually resulted in partial inhibition of tumor cell lysis, but nearly complete inhibition of lysis was observed if all three receptors were blocked simultaneously (5). NKp46 has also been implicated in recognition of virus-infected cells through its capacity to bind to viral hemagglutinins (6-8).

#### References:

1. Moretta, L. and A. Moretta (2004) *EMBO J.* **23**:255.
2. Ferlazzo, G. *et al.* (2004) *J. Immunol.* **172**:1455.
3. Biassoni, R. *et al.* (1999) *Eur. J. Immunol.* **29**:1014.
4. Westgaard, I. *et al.* (2004) *J. Leukoc. Biol.* PMID 15356098.
5. Pende, D. *et al.* (1999) *J. Exp. Med.* **190**:1505.
6. Arnon, T. *et al.* (2004) *Blood* **103**:664.
7. Arnon, T. *et al.* (2001) *Eur. J. Immunol.* **31**:2680.
8. Mandelboim, O. *et al.* (2001) *Nature* **409**:1055.

#### PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.