

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human NKp44/NCR2 in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant mouse MAR-1, recombinant human (rh) NKp30, rhNKp46, or rhNKp80 is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 253422
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human NKp44/NCR2 Gln22-Pro190 Accession # O95944
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

NKp44, along with NKp30 and NKp46, constitute a group of receptors termed "Natural Cytotoxicity Receptors" (NCR) (1). These receptors are expressed almost exclusively by NK cells and play a major role in triggering NK-mediated killing of most tumor cell lines. No rodent ortholog to NKp44 has been identified. Human NKp44, also known as NCR2, is a 44 kDa type I transmembrane glycoprotein that is characterized by the presence of one extracellular V-like immunoglobulin domain (2). It is synthesized as a 276 amino acid (aa) precursor that contains a 21 aa signal sequence, a 171 aa extracellular region, a 21 aa transmembrane segment and a 63 aa cytoplasmic tail. Alternate splicing in both the cytoplasmic tail and extracellular region generates multiple isoforms of unknown significance. The Ig-like region is unaffected. A physical association with the ITAM-bearing accessory protein, DAP12, occurs via a charged residue in the NKp44 transmembrane domain. Ligation of NKp44 with a specific antibody results in phosphorylation of DAP12 (3) and activation of target cell lysis in a redirected killing assay (4). NKp44 is absent from resting NK cells but is upregulated upon activation with IL-2. Activation-induced expression occurs in the CD56^{dim} CD16⁺ NK subset that accounts for more than 85% of NK cells found in peripheral blood and spleen, as well as the CD56^{bright} CD16⁻ NK subset that constitutes the majority of NK cells in lymph node and tonsil (5). Studies with neutralizing antibodies reveal that NKp44 is partially responsible for triggering lytic activity against several tumor cell types (2, 6). Blocking any of the individual NCRs results in partial inhibition of tumor cell lysis, but nearly complete inhibition of lysis is observed if all three receptors are blocked simultaneously (6). NKp44 has also been implicated in recognition of virus-infected cells through its capacity to bind to viral hemagglutinins (7).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.