

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human ATM in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human ATR is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 664703
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human ATM Arg2138-Arg2400 Accession # Q13315
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunocytochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

ATM (Ataxia Telangiectasia Mutated) is a 350-370 kDa member of the ATM subfamily, PI3/PI4-kinase family of enzymes. It is ubiquitously expressed, and serves as a DNA damage sensor. ATM is activated via autophosphorylation at double strand breaks. Following activation, multiple substrates are phosphorylated, including Chk2, and ATR is recruited and activated as part of an integrated repair circuit. Human ATM is 3056 amino acids (aa) in length. It contains one FAT (focal adhesion targeting) domain (aa 1960-2566), a PI-3/PI-4 kinase catalytic domain (aa 2712-2962) and a second, C-terminal FAT domain (aa 3024-3056). There are at least six Ser and four Thr utilized phosphorylation sites, and one critical acetylation activation site at Lys3016. There are at least four potential splice variants. One shows a Trp substitution for aa 536-3056, a second contains an eight aa substitution for aa 2506-3056, a third possesses a five aa substitution for aa 1637-3056, while a fourth contains a premature truncation after Lys2756. Over aa 2138-2400, human ATM shares 82% aa identity with mouse ATM.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.