

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Persephin in Western blots. In Western blots, no cross-reactivity with recombinant human (rh) AgRP, rhArtemin, rhβIG-H3, rhCrip1, recombinant drosophila Decapentaplegic, rhGDNF, rhLAP, rhLatent TGF-β1, rhLefty-A, rhMIS, rhNeurtu
Source	Monoclonal Mouse IgG _{2B} Clone # 275508
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human Persephin Ala61-Gly156 Accession # O60542
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Persephin is a secreted protein belonging to the glial cell line-derived neurotrophic factor (GDNF) family of the TGF-β superfamily. It shares 38-46% amino acid (aa) identity with family members GDNF, neurturin and artemin. Persephin is expressed at very low levels in most tissues (1). The 156 aa, 10-12 kDa mature protein contains a signal sequence, a pro-domain and a 96 aa mature sequence with several cysteines that are conserved among family members. It circulates as an unglycosylated disulfide-linked homodimer. Mature human Persephin shares 81% and 80%, 89%, and 87% amino acid sequence identity with mouse, rat, bovine and canine Persephin, respectively. Like other GDNF family members, Persephin acts through engagement of GRFα4, a glycosylphosphatidylinositol (GPI)-linked GDNF receptor family (GRF) member that signals through the receptor tyrosine kinase RET. Persephin is reported to promote both the survival and growth of central dopaminergic and motor neurons, and kidney development (1). These effects are correlated with the expression patterns of GRFα4, and RET (2, 3). Functional GRFα4 isoforms are found only in thyroid, adrenal medulla and portions of the central nervous system, and include GPI-linked, transmembrane and soluble forms (3, 4). *In vitro*, Persephin promotes survival only in neurons which coexpress GPI-linked GRFα4 with RET (2, 5). This effect does not show a strong correlation to the recruitment of RET in lipid rafts seen with other GDNF family members (6). Disruption of the Persephin gene results in mice that are morphologically normal but have more damage and less effective repair after central nervous system insult that stimulates a stroke. Microinjection of Persephin prior to treatment protects against damage in both wild-type and mutant mouse brains, but surprisingly, high doses of Persephin are detrimental (7).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.