

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human FGF-6 in ELISAs. In direct ELISAs, approximately 75% cross-reactivity with recombinant human (rh) FGF-5 and less than 10% cross-reactivity with rhFGF-4 is observed. Does not neutralize the activity of FGF acidic, FGF basic, rhFGF-4, or
Source	Monoclonal Mouse IgG ₁ Clone # 27028
Purification	Protein A or G purified from ascites
Immunogen	<i>E. coli</i> -derived recombinant human FGF-6 Ala53-Ile208 Accession # P10767
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

ELISA Capture (Matched Antibody Pair)	Optimal dilution of this antibody should be experimentally determined.
ELISA Detection (Matched Antibody Pair)	Optimal dilution of this antibody should be experimentally determined.
Neutralization	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Fibroblast Growth Factor-6 (FGF-6), also known as HST-2, is a 25-28 kDa member of the FGF family of heparin binding polypeptides which are potent regulators of cell proliferation, differentiation, and function. FGF proteins contain a 120 amino acid (aa) core FGF domain that exhibits a β-trefoil structure (1, 2). Mature human FGF-6 is a 171 aa protein that shares 94% aa sequence identity with mouse and rat FGF-6 (3). It binds and signals primarily through FGF R1c, 2c, and 4 (4). FGF-6 functions as a mitogen for fibroblasts, vascular endothelial cells, and prostate carcinoma cells, and N-linked glycosylation is required for the full mitogenic effect (5-7). FGF-6 expression is restricted to skeletal muscle during development although it can be upregulated in prostate cancer and Kaposi sarcoma (7-9). In the adult, FGF-6 is upregulated in injured skeletal muscle and is required for muscle regeneration (10). FGF-6 inhibits the terminal differentiation of myoblasts and also cooperates with TGF-β2 to promote chondrogenesis in embryonic somites (8, 11).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.