

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Serpin G1/C1 Inhibitor in direct ELISAs and Western blots. In Western blots, no cross-reactivity with recombinant human Serpin A1, A3, A4, A5, C1, D1, E2, or F1 is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 350507
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Human plasma-derived Serpin G1/C1 Inhibitor
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunoprecipitation	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

As a member of the serpin superfamily of serine protease inhibitors, Serpin G1/C1 inhibitor is the physiological inhibitor of activated C1r and C1s, two serine proteases involved in the classical complement pathway. In addition, it inhibits plasma kallikrein and coagulation factor XIIa, two serine proteases involved in the processing of kininogen to release bradykinin. Therefore, it plays an important role in regulating activation of both the complement and contact systems (1). Serpin G1 deficiency results in hereditary angioedema, which is characterized by recurrent episodes of localized angioedema of the skin, gastrointestinal mucosa or upper respiratory mucosa (2). The deduced amino acid sequence of human Serpin G1 precursor consists of 500 residues with a signal peptide. The mature protein of 478 amino acid residues is heavily glycosylated (1).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.