

DESCRIPTION

| | |
|---------------------------|--|
| Species Reactivity | Human |
| Specificity | Detects human LIF R α in direct ELISAs and Western blots. |
| Source | Monoclonal Mouse IgG ₁ Clone # 32953 |
| Purification | Protein A or G purified from ascites |
| Immunogen | <i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant human LIF R α Gln45-Ser833 Accession # P42702 |
| Conjugate | Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm |
| Formulation | Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

| | Recommended Concentration | Sample |
|-----------------------|--------------------------------------|--|
| Flow Cytometry | 0.25-1 μ g/10 ⁶ cells | HeLa human cervical epithelial carcinoma cell line |

PREPARATION AND STORAGE

| | |
|--------------------------------|--|
| Shipping | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. |
| Stability & Storage | Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied. |

BACKGROUND

The activities of the pleiotropic cytokine LIF are mediated through a high-affinity heterodimeric receptor complex consisting of two membrane glycoproteins: an α subunit (LIF R α , also known as LIF R and CD118) that binds LIF with low affinity and the 130 kDa (gp130) subunit that does not bind LIF by itself, but is required for high-affinity binding of LIF by the complex. The gp130 subunit was first described as the signal transducing subunit of the high-affinity IL-6 receptor complex. Besides LIF, the high-affinity heterodimeric LIF receptor complex has been shown to mediate the activities of oncostatin M (OSM), cardiotrophin-1 and ciliary neurotrophic factor (CNTF).

Human LIF R α cDNA encodes a 1097 amino acid (aa) residue precursor type I membrane protein with a 44 aa residue signal peptide, a 789 aa residue extracellular domain, a 26 aa residue transmembrane domain, and a 238 aa residue cytoplasmic domain. LIF R α is a member of the cytokine receptor family and has extensive homology to gp130. The extracellular domain of LIF R α has two cytokine receptor domains and three fibronectin type III repeats. In mouse, mRNAs encoding a soluble LIF R α and lacking transmembrane and intracellular domains, have been isolated. Soluble LIF R α has been shown to bind LIF and has LIF antagonistic activity.

References:

1. Bazan, J.F. 1990, Proc. Natl. Acad. Sci. USA **87**:6934.
2. Gearing, D.P. (1994) *Guidebook to Cytokines and Their Receptors*, Academic Press, p130.
3. Pennica D. *et al.* (1995) J. Biol. Chem. **270**:10915.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.