

## DESCRIPTION

<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse Chordin-like 2/CHRD2 in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant human CHRD2, recombinant mouse (rm) CHRD1, or rmChordin is observed.
<b>Source</b>	Monoclonal Rat IgG <sub>2A</sub> Clone # 328905
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant mouse Chordin-like 2/CHRD2 Gln24-Leu426 Accession # Q8VEA6.1
<b>Conjugate</b>	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
<b>Formulation</b>	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

<b>Western Blot</b>	Optimal dilution of this antibody should be experimentally determined.
<b>Immunohistochemistry</b>	Optimal dilution of this antibody should be experimentally determined.
<b>Immunoprecipitation</b>	Optimal dilution of this antibody should be experimentally determined.

## PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

## BACKGROUND

Chordin-like 2 (CHL2; also known as breast tumor novel factor 1/BNF-1), is a secreted glycoprotein that has significant homology to chordin and acts as a TGF-β superfamily antagonist (1-3). CHL2 is a chordin family member with structural homology to CHL1 (also known as neuralin or ventroptin) (1, 2). Mouse CHL2<sub>401</sub> cDNA encodes a 45-50 kDa, 426 amino acid (aa) residue precursor protein with a putative 25 aa signal peptide and a 401 aa mature segment. The mature segment contains three 63 aa cysteine-rich von Willebrand type C repeats (CRs) that are conserved among chordin family members in the spacing of 10 cysteine residues (1, 2). It also contains two potential N-linked glycosylation sites and one putative NLS that lies just proximal to the third CR repeat.

Mouse CHL2, like human CHL2, appears to undergo extensive alternate splicing. This splicing generates both secreted and intracellular forms of CHL2, and influences the type of TGF-β superfamily member bound (1, 2). The CRs of chordin, especially CR1 and CR3, have been shown to be the functional domains for BMP binding (4). The CR1 and CR3 of CHL proteins are most similar to CR3 of chordin (1). Mature mouse and human CHL2 share 71% amino acid identity, while mouse CHL1 and -2 share 40% amino acid identity (2). Like chordin, CHL2<sub>401</sub> exhibits BMP inhibitory activity by directly interacting with BMP-4 and preventing binding to its receptor (1). However, another CHL2 isoform with an additional exon 9b has been shown to bind and inhibit Activin A activity as well (2). CHL2<sub>401</sub> is not abundantly expressed in mouse embryos, but is detected only in the chondrocytes of developing joints and in the connective tissue of reproductive organs (1). Mouse CHL2 acts to reduce the rate of matrix accumulation in mesenchymal cells, acting as a negative regulator of cartilage formation (1). In the adult mouse, CHL2 is again detected only faintly in liver, kidney, skeletal muscle and testis (1). Expression patterns in human tissue blots are distinct from those expressed in mouse (1, 2).

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.