

# Human IL-5 R $\alpha$ /CD125 Alexa Fluor® 488-conjugated Antibody

Monoclonal Mouse IgG<sub>1</sub> Clone # 26815

Catalog Number: FAB253G

100  $\mu$ g

## DESCRIPTION

<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human IL-5 R $\alpha$ /CD125 in direct ELISAs and Western blots. In direct ELISAs and Western blots, this antibody does not cross-react with recombinant human (rh) IL-5 R $\beta$ , rhIL-9 R, rhIL-4 R, rhIL-5 R $\alpha$ , or rhIL-13 R $\alpha$ 1.
<b>Source</b>	Monoclonal Mouse IgG <sub>1</sub> Clone # 26815
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant human IL-5 R $\alpha$ /CD125 Asp21-Arg335
<b>Conjugate</b>	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
<b>Formulation</b>	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Flow Cytometry</b>	0.25-1 $\mu$ g/10 <sup>6</sup> cells	HEK293 human embryonic kidney cell line transfected with human IL-5 R $\alpha$ /CD125 and eGFP

## PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Protect from light. Do not freeze.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul>

## BACKGROUND

Interleukin 5, produced primarily by activated T cells and mast cells, has diverse biological effects on a variety of cell types. Human IL-5 is a potent eosinophil differentiation and activation factor *in vivo* and *in vitro*. Additionally, it has also been reported that IL-5 can stimulate the proliferation and/or differentiation of basophils and B cells. The multiple effects of IL-5 are mediated by binding of the cytokine to specific cell surface receptors expressed on target cells. As is the case with many other cytokines, the functional high-affinity receptor for IL-5 is a complex consisting of a ligand binding subunit ( $\alpha$  chain) and a second subunit ( $\beta$  chain) that can modulate the ligand binding affinity of the receptor complex. In the case of IL-5, the  $\beta$  subunit is shared with the high affinity receptor complexes for IL-3 and GM-CSF. The  $\beta$  chain does not bind any of the cytokines in question but is indispensable for the cytokine-mediated signaling. cDNA clones for the  $\alpha$  chain (IL-5 R $\alpha$ ) of both the mouse and human high affinity IL-5 receptor complexes have been isolated. Human and mouse IL-5 R $\alpha$  are both members of the hematopoietin receptor superfamily characterized by the presence of the WSXWS, and a four cysteine residue motif in the extracellular domain of the transmembrane protein. In addition to the cDNA clone encoding the full-length transmembrane protein, cDNA clones that arise from alternative splicing and that encode soluble secreted forms of IL-5 R $\alpha$  have been isolated from mouse as well as human cells. A naturally-occurring soluble form of the IL-5 R $\alpha$  has been detected in biological fluids of autoimmune-prone mice and mice bearing chronic B cell leukemia (BCL<sub>1</sub>). A recombinant human IL-5 soluble receptor  $\alpha$  has been shown to bind the human IL-5 dimer in a 1:1 ratio and acts as a human IL-5 antagonist. This molecule inhibits the proliferation of IL-5-dependent cell lines and blocks human umbilical cord blood eosinophil differentiation.

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.