

DESCRIPTION


| | |
|---------------------------|--|
| Species Reactivity | Human |
| Specificity | Detects human NG2/MCSP in Western blots. |
| Source | Monoclonal Mouse IgG ₁ Clone # LHM-2 |
| Purification | Protein A or G purified from hybridoma culture supernatant |
| Immunogen | Human melanoma cells |
| Conjugate | PerCP (Peridinin-chlorophyll Protein Complex) Excitation Wavelength: 482 and 564 nm Emission Wavelength: 675 nm |
| Formulation | Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

| | Recommended Concentration | Sample |
|-----------------------|----------------------------------|---------------|
| Flow Cytometry | 10 μ L/10 ⁶ cells | See Below |

DATA



PREPARATION AND STORAGE

| | |
|--------------------------------|---|
| Shipping | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. |
| Stability & Storage | Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied. |

BACKGROUND

NG2, also known as MCSP, is a chondroitin sulfate proteoglycan that has been used as a cell surface marker for melanoma (1) and glial precursor cells (2). NG2 also promotes epidermal stem cell patterned clustering (3).

References:

1. Kupsch, J.M. *et al.* (1995) *Melanoma Res.* **5**:403.
2. Stegmuller, J. *et al.* (2002) *J. Neurocytol.* **31**:497.
3. Legg, J. *et al.* (2003) *Development* **130**:6049.