

DESCRIPTION


Species Reactivity	Mouse
Specificity	Detects mouse ULBP-1 in flow cytometry.
Source	Monoclonal Rat IgG _{2A} Clone # 237104
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	BaF3 mouse pro-B cell line transfected with mouse ULBP-1
Conjugate	PerCP (Peridinin-chlorophyll Protein Complex) Excitation Wavelength: 482 and 564 nm Emission Wavelength: 675 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage **Protect from light. Do not freeze.**

- 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

ULBP-1, also known as MULT-1 (Mouse UL16-Binding Protein-like Transcript 1), is a 53 kDa, MHC Class I-like molecule that belongs to the mouse family of NKG2D ligands (1-4). It is a type I transmembrane glycoprotein that is synthesized as a 334 amino acid (aa) precursor. It contains a 25 aa signal sequence, a 186 aa extracellular region, a 19 aa transmembrane segment and a 104 aa cytoplasmic tail (2). The extracellular region contains an α 1 and α 2 like domain with two intrachain disulfide bonds. ULBP-1 is distantly related to other human and mouse NKG2D ligands, and more distantly related to the MHC class I proteins (3). Unlike most NKG2D ligands, transcripts for ULBP-1 have been detected in a wide variety of mouse tissues and tumor cells lines (3). The receptor for ULBP-1 is NKG2D, a 35 kDa C-type lectin that is found on mouse NK cells, activated CD8⁺ T cells, epidermal $\gamma\delta$ T cells, and activated macrophages (1, 5, 6, 7). Recombinant ULBP-1 protein binds to NKG2D with high affinity ($K_D = 6$ nM) (2). Although an activating receptor, general cellular responses to NKG2D ligation depend upon the isoform of NKG2D and the cell type (5). Exposure to immobilized ULBP-1 or ULBP-1-transfected cells elicits IFN- γ production by NK cells (3). Ectopic expression of ULBP-1 on the RMA mouse tumor cell line leads to tumor rejection in syngeneic mice (3).

References:

1. Raulet, D.H. (2003) *Nat. Rev. Immunol.* **3**:781.
2. Carayannopoulos, L. *et al.* (2002) *J. Immunol.* **169**:4079.
3. Diefenbach, A. *et al.* (2003) *Eur. J. Immunol.* **33**:381.
4. Krmpotic, A. *et al.* (2005) *J. Exp. Med.* **201**:211.
5. Diefenbach, A. *et al.* (2002) *Nat. Immunol.* **3**:1142.
6. Ho, E.L. *et al.* (1998) *Proc. Natl. Acad. Sci. USA* **95**:6320.
7. Carayannopoulos, L.N. *et al.* (2002) *Eur. J. Immunol.* **32**:597.