

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Cardiotrophin-1/CT-1 in direct ELISAs and Western blots.
Source	Monoclonal Mouse IgG ₁ Clone # 89215
Purification	Protein A or G purified from ascites
Immunogen	<i>E. coli</i> -derived recombinant human Cardiotrophin-1/CT-1 Ser2-Ala201 Accession # Q16619
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Cardiotrophin-1 (CT-1) is a member of the cytokine family which also includes IL-6, IL-11, leukemia inhibitory factor (LIF), oncostatin M (OSM), and ciliary neurotrophic factor (CNTF). It was originally isolated based on its ability to induce cardiac myocyte hypertrophy *in vitro*. CT-1 has since been shown to be a pleiotrophic cytokine with overlapping actions with other IL-6 family members on a variety of cell types. Human CT-1 encodes a 201 amino acid (aa) residue protein that lacks a hydrophobic signal peptide. The mechanism of CT-1 release from cells is currently not understood. Human and mouse CT-1 share 80% aa sequence identity and exhibit cross-species activity. CT-1 is highly expressed in heart, skeletal muscle, liver, lung and kidney. Lower levels of CT-1 expression is also seen in testis and brain. CT-1 initiates downstream signaling pathways through the heterodimerization of gp130 and the LIF receptor β subunit. A third α receptor subunit has been implicated in the receptor complex.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.