

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Betacellulin/BTC in direct ELISAs and Western blots. In direct ELISAs, no cross-reactivity with recombinant mouse (rm) BTC, recombinant human (rh) Amphiregulin, rhEGF, rhHB-EGF, rhHeregulin-α, rhHeregulin-β, rhTGF-α, or rmEpiregulin is
Source	Monoclonal Mouse IgG ₁ Clone # 34301
Purification	Protein A or G purified from ascites
Immunogen	<i>E. coli</i> -derived recombinant human Betacellulin/BTC Asp32-Tyr111 Accession # P35070
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Betacellulin (BTC) is a new member of the EGF family of cytokines that is comprised of at least ten proteins including EGF, TGF-α, amphiregulin, HB-EGF, and the various heregulins. All of these cytokines are synthesized as transmembrane precursors and are characterized by the presence of one or more EGF structural units in their extracellular domain. The soluble forms of these cytokines are released by proteolytic cleavage. BTC, a heparin-binding protein, was originally isolated from the conditioned media of mouse pancreatic beta tumor cells as a 32 kDa glycoprotein composed of 80 amino acid residues. The cDNA encoding human BTC was cloned from a human breast adenocarcinoma cell line (MCF-7) cDNA library. Human and mouse cDNAs encode BTC precursor proteins of 178 and 177 amino acid residues, respectively. At the amino acid sequence level, human BTC precursor protein exhibits 79% identity with that of the mouse BTC precursor. In a mouse cell line transfected with human BTC cDNA, three forms of soluble human BTC have been detected: the glycosylated, intact BTC composed of 80 amino acid residues, a truncated molecule lacking 12 amino acid residues from the amino terminus, and a second truncated molecule lacking 30 amino acid residues from the amino terminus. The biological activities of these BTC forms were shown to be identical. BTC can bind to the EGF receptor and is a potent mitogen for Balb/c 3T3 fibroblasts, retinal pigment epithelial cells and vascular smooth muscle cells.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.