

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Kallikrein 14 in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 337903
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Kallikrein 14 Gln19-Met248 Accession # AAD50773
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Immunoprecipitation	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Human tissue kallikreins refer to a group of secreted serine proteases that are encoded by homologous genes clustering on chromosome 19q13.3-4. As a member of this family, human tissue Kallikrein 14 (hKLK14) is present in many tissues, with high levels in breast, skin, prostate, and brain. The 251 amino acid hKLK14 precursor consists of a signal peptide (residues 1 to 18), a pro peptide (residues 19 to 24) and an active protein (residues 25 to 251) (1). Its enzymatic activity has been shown to be mainly trypsin-like (2). However, its physiological substrates and functions are still unclear. Several studies have suggested that hKLK14 may have clinical utility as a biomarker for cancer of the breast, ovary, and prostate (3, 4). In addition, it may be the initiator of a kallikrein proteolytic cascade responsible for the degradation of the adhesion structures in the stratum corneum (2). The purified, secreted rhKLK14 corresponds to the pro form (residues 19 to 248) with a replacement of the last three residues with a His tag at the C-terminus. When activated by thermolysin, it displays enzymatic activity towards a fluorogenic peptide described above. This activity can be inhibited by rhSerpin A4, E1, and F2 (R&D Systems, Catalog # 1669-PI, 1786-PI, and 1470-PI, respectively).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.