

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human NT-4 in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant human (rh) BDNF or rhNT-3 is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 36507
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human NT-4 Gly81-Ala210 (predicted) Accession # P34130
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Neurotrophin-4 (NT-4), also known as NT-5, is a member of the NGF family of neuronal and epithelial growth factors. Neurotrophins have six conserved cysteine residues that are involved in the formation of three disulfide bonds (1-3). The human NT-4 cDNA encodes a 210 amino acid (aa) precursor that includes a 24 aa signal sequence, a 56 aa propeptide, and a 130 aa mature protein (4, 5). NT-4 is synthesized as a 28 kDa prepropeptide that is proteolytically processed to generate the mature protein. Mature human NT-4 shares 48-52% aa sequence identity with human beta-NGF, BDNF, and NT-3. It shares 91% and 95% aa sequence identity with mouse and rat NT-4/5, respectively. The mature protein is secreted as a homodimer and can also form heterodimers with BDNF or NT-3 (6). NT-4 binds and induces receptor dimerization and activation of TrkB (4, 7). NT-4 promotes the development and survival of selected peripheral and CNS neurons (8-10). BDNF, which also activates TrkB, overlaps with many but not all NT-4 functions, a distinction that is likely due to differences in expression patterns (8-10). NT-4 induced TrkB signaling augments NMDA receptor activity and increases neuronal sensitivity to excitotoxic cell death (11). It also promotes the proliferation of keratinocytes and accelerates hair follicle regression during the follicular cycle (12, 13). NT-4 is secreted by activated T cells and granulocytes at sites of inflammation where it contributes to tissue regeneration (14-16).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.