

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CL-P1/COLEC12 in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant mouse CL-P1 is observed.
Source	Monoclonal Rat IgG ₁ Clone # 388305
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human CL-P1/COLEC12 isoform 1 Leu57-Leu742 Accession # Q5KU26
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Collectins are a family of Ca⁺⁺-dependent, C-type lectins that contain a collagenous domain and function as recognition molecules for molecular patterns found on pathogens (1-4). Human collectin placenta 1 (CL-P1; also known as collectin sub-family member 12 and SRCL type I [scavenger receptor with C-type lectin type II]) is a 110 kDa member of the collectin family of glycoproteins (5, 6). With two exceptions, all collectins are secreted. CL-P1 is the only collectin known to be membrane bound, while CL-L1 (collectin liver-1) is the only known cytoplasmic collectin (1). Human CL-P1 is synthesized as a 742 amino acid (aa) type II transmembrane glycoprotein that contains an N-terminal 39 aa cytoplasmic domain, a 17 aa transmembrane segment, and a 686 aa C-terminal extracellular region (6). The short cytoplasmic domain contains an internalization motif (Y-K-R-F) while the extracellular region is complex, demonstrating a coiled-coil segment, a Ser-Thr rich region, a collagen-like structure and a C-type lectin/ carbohydrate recognition domain (CRD). Notably, this CRD recognizes galactose (and fucose) within the context of asialo-orosomucoids associated with the Lewis^x epitope (7, 8). CL-P1 has a 300 kDa trimeric form due to its collagen-like and coiled-coil helical domains (1, 5). There is a 97 kDa, alternate splice form of CL-P1 (SRCL type II) that shows a 120 aa truncation at the C-terminus. This effectively removes the entire CRD found on full-length CL-P1 (6). Human CL-P1 shares 93% aa sequence identity with mouse CL-P1 over the entire extracellular region, and 87% aa identity within each species CRD (5, 9). Human CL-P1 is expressed in vascular endothelial cells (5). CL-P1 may play a role in bacterial recognition or as a scavenger receptor for desialylated glycoproteins.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.