

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human MFG-E8 in direct ELISAs and Western blots. In direct ELISAs and Western blots, this antibody does not cross-react with recombinant human (rh) C1r, rhC1s, rhC2, or rmMFG-E8.
Source	Monoclonal Mouse IgG _{2A} Clone # 278901
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human MFG-E8 Leu24-Cys387 Accession # Q08431
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Milk Fat Globulin Protein E8 (MFG-E8), also known as Lactadherin, MP47, breast epithelial antigen BA46, and SED1, is a 66-75 kDa pleiotropic secreted glycoprotein that promotes mammary gland morphogenesis, angiogenesis, and tumor progression. MFG-E8 also plays an important role in tissue homeostasis and the prevention of inflammation (1). Human MFG-E8 contains one N-terminal EGF-like domain and two C-terminal F5/8-type discoidin-like domains (2). It shares 63% and 61% aa sequence identity with comparable regions of mouse and rat MFG-E8, respectively. Shorter isoforms of human MFG-E8 may have N-terminal deletions (beginning near the end of the first discoidin-like domain), internal deletions (lacking either the EGF-like domain or the central region of the second discoidin-like domain), or C-terminal deletions (truncated within the second discoidin-like domain) (3). A 50 aa internal proteolytic fragment of human MFG-E8 (known as Medin) is a major component of aortic medial amyloid deposits (4). MFG-E8 is released into the milk in complex with lipid-containing milk fat globules. It is also found in multiple other cell types including endothelial cells and smooth muscle cells of the vasculature, immature dendritic cells, at the acrosomal cap of testicular and epididymal sperm, and in epithelial cells of the endometrium (1). MFG-E8 binds to the Integrins $\alpha V\beta 3$ and $\alpha V\beta 5$ and potentiates the angiogenic action of VEGF through VEGF R2 (5, 6). It reduces inflammation and tissue damage in a variety of settings. MFG-E8 functions as a bridge between phosphatidylserine on apoptotic cells and Integrin $\alpha V\beta 3$ on phagocytes, leading to the clearance of apoptotic debris (7). It mediates the engulfment of apoptotic bodies in atherosclerotic plaques and prion-infected brain (8, 9) and of apoptotic B cells during germinal center reactions (10, 11). MFG-E8 also promotes the removal of excess Collagen in fibrotic lungs and the regeneration of damaged intestinal epithelia (12, 13). Its tissue-protective role impairs anti-tumor immunity and chemotherapy-induced apoptosis (14). MFG-E8 in the breastmilk blocks rotavirus infection in nursing babies (15).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.