

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human FLRT2 in direct ELISAs and Western blots. In direct ELISAs and Western blots, this antibody does not cross-react with recombinant human (rh) FLRT1 or rhFLRT3.
Source	Monoclonal Mouse IgG _{2B} Clone # 367205
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human FLRT2 Cys36-Ser539 Accession # O43155
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

FLRT2 is one of three FLRT (fibronectin, leucine rich repeat, transmembrane) glycoproteins expressed in distinct areas of the developing brain and other tissues (1, 2). The 85 kDa type I transmembrane (TM) human FLRT2 is synthesized as a 660 amino acid (aa) precursor with a 35 aa signal sequence, a 506 aa extracellular domain (ECD), a 21 aa TM segment and a 98 aa cytoplasmic region. The ECD contains 10 N-terminal leucine-rich repeats flanked by cysteine-rich areas, and a juxtamembrane fibronectin type III domain (1). The human FLRT2 ECD shares 97%, 96%, 99%, 96% and 95% aa sequence identity with mouse, rat, equine, canine and bovine FLRT2 ECD, respectively. Human FLRT1 and FLRT3 ECDs share approximately 47% aa identity with FLRT2. The fibronectin domain of all three FLRTs can bind to FGF receptors (2). This binding is thought to regulate FGF signaling during development (2, 3). The LRR domains are responsible for both the localization of FLRTs in areas of cell contact and homotypic cell-cell association (4). This may be through direct interactions with other FLRT molecules or, as has been shown for FLRT3, by regulating internalization of adhesion molecules such as cadherins (4, 5). In adulthood, FLRT2 mRNA is most abundant in pancreas, but is also present in skeletal muscle, brain and heart (1). FLRT2 in mouse embryos shows highest expression in a subset of the sclerotome in the brain, the stomach, and posterior to the developing heart (2). This expression is distinct from that of FLRT1 and FLRT3 (2).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.