

DESCRIPTION

Species Reactivity	Human/Mouse
Specificity	Detects the pro form of human or mouse Meprin β Subunit/MEP1B in Western blots. It does not react with the mature form.
Source	Monoclonal Rat IgG ₁ Clone # 289731
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Human embryonic kidney cell line HEK293-derived recombinant human Meprin β Subunit/MEP1B Thr23-Ser293 Accession # Q16820
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Meprins are multimeric proteases composed of α and β subunits, which are members of the astacin family of zinc endopeptidases (1, 2). Both subunits form disulfide-linked homo- or heterooligomers, which are also referred to as Meprin A (composed of α subunits with or without β subunits) and Meprin B (composed of β subunits only) (3). Although the two subunits share 42% amino acid (aa) sequence identity, they differ significantly in their oligomeric structure, post-translational processing and subsequently cellular location, and substrate and peptide bond specificity (4). The 701 aa sequence of human Meprin β subunit precursor consists of a signal peptide (aa 1 to 21), a pro region (aa 22 to 61), and a mature chain (aa 62 to 701) containing catalytic (aa 62 to 259), MAM (aa 260 to 429), MATH (aa 430 to 585), EGF-like (aa 604 to 644), transmembrane (aa 653 to 673), and cytoplasmic (aa 674 to 701) domains.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.