

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human BMP-15/GDF-9B in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant human BMP-1.1, -2, -3, -3b, -4, -5, -6, -7, -8, or -10 is observed.
Source	Monoclonal Rat IgG _{2A} Clone # 317809
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human BMP-15/GDF-9B Gln268-Arg392 Accession # O95972
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Bone morphogenetic protein 15 (BMP-15), also known as GDF-9B, is a TGF- β superfamily ligand that is expressed by oocytes throughout folliculogenesis, and plays an important role in oocyte development (1). BMP-15 promotes the FSH-independent proliferation of ovarian granulosa cells (GC) and induces GC glycolysis and cholesterol synthesis (2-4). It also induces GC production of stem cell factor which, in turn, negatively regulates BMP-15 expression in oocytes (5). BMP-15 blocks the FSH-induced GC expression of FSH R and multiple steroidogenic molecules (6). BMP-15 is synthesized with a 249 amino acid (aa) N-terminal propeptide (7). The propeptide is cleaved intracellularly from the 50 kDa proBMP-15 but remains associated with mature BMP-15 (8). Mature BMP-15 exists in 16 kDa and 17 kDa forms which are distinguishable by the presence of O-linked glycosylation on the 17 kDa form (8). Mature BMP-15 is phosphorylated, a modification which is required for the stimulation of GC proliferation (9). BMP-15 exerts its effects through interactions with BMPR-IB/ALK6 and BMPR-II (9-11). Mature BMP-15 forms 34 kDa noncovalently-linked homodimers and 37 kDa heterodimers with mature GDF-9 (12). Both of these oocyte-expressed factors lack the cysteine that mediates disulfide-linked dimerization in most TGF- β superfamily ligands (1). Although heterodimerization with GDF-9 may limit the secretion of active BMP-15, these two factors synergize in promoting oocyte survival and folliculogenesis (12, 13). Mature human BMP-15 shares 70%, 68%, and 78% aa sequence identity with mouse, rat, and sheep BMP-15, respectively. It shares 27%-38% aa sequence with other BMPs.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.