

Human MICL/CLEC12A Alexa Fluor® 700-conjugated Antibody

Recombinant Monoclonal Rabbit IgG Clone # 2439B Catalog Number: FAB29461N

100 µg

DESCRIPTION			
Species Reactivity	Human		
Specificity	Detects human MICL/CLEC12A in direct ELISAs.		
Source	Recombinant Monoclonal Rabbit IgG Clone # 2439B		
Purification	Protein A or G purified from cell culture supernatant		
Immunogen	Mouse myeloma cell line NS0-derived recombinant human MICL/CLEC12A Thr67-Ala265 Accession # Q5QGZ9		
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm		
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.		
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.		

APPLICATIONS				
Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.				
	Recommended	Sample		
	Concentration			
Flow Cytometry	0.25-1 μg/10 ⁶ cells	Human peripheral blood granulocytes		

PREPARATION AND STORAGE		
Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.	
Stability & Storage	s Storage Protect from light. Do not freeze.	
	12 months from date of receipt, 2 to 8 °C as supplied.	

BACKGROUND

Human MICL (myeloid inhibitory C-type lectin-like receptor; also CLL-1 and KLRL-1) is a 75 kDa member of the group V C-type lectin-like receptor family of molecules. Group V molecules have a lectin-type domain that binds non-sugar ligands. MICL is a 265 amino acid (aa) type II transmembrane (TM) glycoprotein that contains a 200 aa extracellular domain (ECD). Multiple isoforms of MICL are reported. One potentially utilizes an alternate start site that adds 10 aa to the cytoplasmic domain. There is one potential soluble form that lacks the TM segment, and two truncated ECD isoforms. Human MICL ECD is 53%, 65%, and 55% aa identical to the ECD in mouse, dog and cow, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

