

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Mimecan in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant human Mimecan is observed.
Source	Monoclonal Rat IgG _{2A} Clone # 329939
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse Mimecan Ala20-Phe298 Accession # Q62000
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Mimecan is a 34 kDa secreted monomeric glycoprotein that belongs to the SLRP, or small leucine-rich proteoglycan family of matrix molecules (1, 2). All SLRP family members contain multiple leucine rich repeats, of which there are two types; a 21 amino acid (aa) proline-containing S-type, and a 26 aa phenylalanine-containing T-type (2, 3). Mimecan is a class III SLRP subfamily member, which means it contains an S-T (vs. S-T-T) repeating motif. Mouse Mimecan is synthesized as a 298 aa precursor that contains a 19 aa signal sequence and a 279 aa mature region (3, 4). The mature region contains a 75 aa N-terminus followed by a 13 aa cysteine-rich region and seven consecutive LRR repeats. There is one (presumably occupied) C-terminal N-linked glycosylation site and a potential keratan sulfate attachment point in LRR #4 (aa 161-180) (3). Nomenclature for the Mimecan molecule is confusing. Mimecan is best thought of as being the full-length, 279 aa glycoprotein (5). Proteolytic removal of the first 56 aa by BMP-1/Tolloid-like proteinases generates the 25 kDa KSPG form (aa 57-279) (3). The 17 kDa, 56 aa mature N-terminus has been referred to as the minecan prosegment (3). When the prosegment is removed, mature mouse KSPG (aa 57-279) is 98%, 91%, 90% and 91% aa identical to rat, bovine, human and canine KSPG, respectively. A 12 kDa form (aa 175-279) consisting of the 105 aa of the C-terminus (OIF/osteoglycan) has also been observed (6). Noncoding alternate splice forms for mouse are believed to exist and should be expressed in a tissue specific fashion (7, 8). No splice forms are known that impact the coding region. Mimecan seems to exist as proteoglycan (PG) and non-PG forms. In cornea, keratan sulfate is present, although the exact function of the PG is unknown. The corneal extracellular matrix is normally transparent. While Mimecan is reported to inhibit fibrillogenesis, possibly by limiting lateral accretion of collagen monomers, it is unclear if the mimican PG actually contributes to transparency (3, 9). In other tissues, Mimecan and its derivatives can exist as simple glycoproteins, absent of any PG adduct (6). The uncertain nature of Mimecan's structure and function gives rise to its name: *Mime* is a Norse-legend dwarf that was notoriously deceitful and misleading (6).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.