

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse LRP-6 in direct ELISAs and Western blots. In direct ELISAs, this antibody shows approximately 50 - 60% cross-reactivity with recombinant human (rh) LRP-6 and no cross-reactivity with rhLDL R.
Source	Monoclonal Rat IgG _{2B} Clone # 290913
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant mouse LRP-6 Met1-Pro1368 Accession # O88572
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

The low-density lipoprotein (LDL) receptor-related protein 5 (LRP-5) and LRP-6 constitute a distinct subgroup of the LDL receptor family (1). Both LRP-5 and LRP-6 are type I transmembrane proteins that function as Wnt co-receptors with Frizzled proteins (FZD) (2, 3, 4). The mouse LRP-6 cDNA encodes a 1613 amino acid (aa) residue precursor including a 19 aa signal sequence, 1351 aa extracellular domain (ECD), a 23 aa transmembrane (TM) segment, and a 20 aa cytoplasmic domain (5). The ECD contains 20 YWTD motif-containing LDLR-B domains, four EGF-like repeats, and three cysteine-rich LDLR-A repeats. The ECD of mouse LRP-6 shares 71% aa sequence identity with the ECD of mouse LRP-5 and 98% aa sequence identity with the ECD of human, rat, and canine LRP-6. The intracellular region of LRP-6 contains repeated PPPSP motifs. When the Ser/Thr in these motifs are phosphorylated, LRP-6 can interact with Axin and propagate canonical Wnt signal transduction (6, 7). LRP-6 forms inactive homodimers via its YWTD-EGF domains (8). Wnt binding to FZD and subsequent association with LRP-6 lead to activating conformational changes in LRP-6 cytoplasmic domains (8). LRP-6 can also interact directly with the Dickkopf (Dkk), sclerostin, and Wise proteins which are modulators of Wnt signaling (9 - 14). Formation of a ternary complex of LRP-6, Dkk-1, and Kremen triggers the internalization of the complex and removal LRP-6 from the cell surface. LRP-5 and LRP-6 share overlapping functions in diverse embryonic developmental processes (15).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.