

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human TGF- β RI/ALK-5 in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 854621R
Purification	Protein A or G purified from cell culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human TGF- β RI/ALK-5 Leu34-Glu125 Accession # P36897
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

ELISA Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

TGF- β RI, also called ALK-5, is an approximately 55 kDa type I transmembrane serine/threonine receptor kinase (1, 2). It contains a cysteine-rich extracellular domain (ECD), a transmembrane helix, and a C-terminal cytoplasmic kinase domain (3). Within the cytoplasmic domain there is also a short, conserved regulatory sequence known as the GS region that is N-terminal to the kinase domain (1). Within the ECD, human TGF- β RI shares 90% and 88% aa sequence identity with mouse and rat TGF- β RI, respectively. In the presence of TGF- β , TGF- β RI forms a complex with, and is phosphorylated by, TGF- β RII (1). Phosphorylated TGF- β RI can then transiently bind and phosphorylate Smad2 and Smad3 (2, 4-6). These phosphorylated Smads form heteromeric complexes with Smad4, translocate to the nucleus, and regulate target gene transcription (2, 4-6). TGF- β RI is likely important during development, since mice deficient for TGF- β RI die at midgestation with severe defects in vascular development of the yolk sac and placenta, and an absence of circulating red blood cells (7). Furthermore, TGF- β RI appears to be involved in proper lymphatic network development (8). Mutations in TGF- β RI have been identified in pancreatic, colorectal, ovarian, and head and neck cancers (9).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.