

## Human NF-M Alexa Fluor® 488-conjugated Antibody

Monoclonal Mouse IgG<sub>1</sub> Clone # 327921

Catalog Number: FAB3029G

DESCRIPTION	
Species Reactivity	Human
Specificity	Detects human NF-M in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant human (rh) NF-H or rhNF-L is observed.
Source	Monoclonal Mouse IgG <sub>1</sub> Clone # 327921
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	E. coli-derived recombinant human NF-M Ser2-Ser358 Accession # P07197
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Shee (SDS) for additional information and handling instructions.

APPLICATIONS		
Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.		
Knockout Validated	Optimal dilution of this antibody should be experimentally determined.	
Western Blot	Optimal dilution of this antibody should be experimentally determined.	

PREPARATION AND STORAGE		
Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.	
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied	

## **BACKGROUND**

Human NF-M (neurofilament-medium length) is a 145-150 kDa phosphorylated glycoprotein that belongs to the type IV intermediate filament family. It is 915 amino acids in length and contains a 100 amino acid (aa) N-terminal globular region, a 320 aa α-helical rod, or central region, and a 500 aa globular tail. The rod region contains a series of heptad repeats that contribute to a coiled-coil interaction with an adjacent NF-M molecule, the first step in intermediate filament oligomerization. The C-terminus shows phosphorylation and O-linked glycosylation. Human NF-M aa 2-358, which includes most of the rod region, shares ~98% aa sequence identity with corresponding region of mouse, rat and canine NF-M.

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

Rev. 9/21/2025 Page 1 of 1

China | info.cn@bio-techne.com TEL: 400.821.3475