

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human FGF-17 in direct ELISAs and Western blots. In direct ELISA, approximately 7% cross-reactivity with recombinant human (rh) FGF-5 is observed and no cross-reactivity with rhFGF-acidic, rhFGF-basic, rhFGF-4, rhFGF-6, rhFGF-7, rhF
Source	Monoclonal Mouse IgG _{2B} Clone # 121630
Purification	Protein A or G purified from ascites
Immunogen	<i>E. coli</i> -derived recombinant human FGF-17 Thr23-Thr216 Accession # O60258
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Fibroblast growth factors (FGFs) play multiple biological functions including angiogenesis, mitogenesis, cellular differentiation and wound repairing. All members of the FGF family have a conserved approximately 120 amino acid core with 30-70% identity. Among FGF family members, FGF-17 is most similar to FGF-8 (60% sequence identity) and FGF-18 (50% sequence identity). The mRNA of FGF-17 was found in midgestation of embryo and multiple adult tissues, and is preferentially expressed in specific sites, such as embryonic brain, developing skeleton and arteries. Human FGF-17 shares 98.6% amino acid (aa) sequence identity with mouse FGF-17. Rat FGF-17 shares 100% aa sequence identity with mouse FGF-17.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.