

DESCRIPTION


Species Reactivity	Rat
Specificity	Detects rat MCAM/CD146 in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant human (rh) MCAM, rhBCAM, or rhALCAM is observed.
Source	Monoclonal Mouse IgG _{2A} Clone # 404722
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant rat MCAM/CD146 Arg19-Lys560 Accession # Q9EPF2
Conjugate	Fluorescein Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm (FITC)
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Melanoma Cell Adhesion Molecule (MCAM), also known as CD146 and MUC18, is a putative Ig-superfamily adhesion molecule that is expressed on endothelial cells and a variety of tumor cells. MCAM is associated with tumor progression and metastasis and may be involved in embryonic neural development. An alternately spliced isoform of rat MCAM has a 40 residue deletion in the cytoplasmic domain. Within the extracellular region, rat MCAM shares 72% and 90% amino acid sequence identity with human and mouse MCAM, respectively.