

DESCRIPTION

Species Reactivity	Human/Mouse
Specificity	Detects human and mouse cIAP-1 and cIAP-2 in Western blots.
Source	Monoclonal Mouse IgG _{2A} Clone # 315301
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human cIAP-2 Asn2-Ser604 Accession # Q13489
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Cellular inhibitor of apoptosis protein1 (cIAP-1, also known as BIRC2, MIHB, and HIAP2) and 2 (cIAP-2, also known as BIRC3, MIHC, and HIAP1), are members of the inhibitor of apoptosis (IAP) family of proteins that inhibit the proteolytic activity of mature caspases. Structurally, cIAP-1 and cIAP-2 are each comprised of 3 BIR (baculovirus inhibitor of apoptosis) domains, a RING finger domain, and a caspase recruitment domain (CARD). Human cIAP-1 and cIAP-2 share 70% sequence identity. Functionally, cIAPs inhibit caspases through the direct interaction of its BIR domain with the active caspase. The ring finger domain of cIAP-1 and -2 also functions as an E3 ubiquitin ligase to ubiquitinate specific target proteins. Caspase activity may be restored by mitochondrial proteins, such as SMAC/Diablo or HtrA2/Omi, through interactions with the Reaper-like motif and the BIR domain. cIAPs are reported to be cleaved by HtrA2/Omi.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.