

DESCRIPTION

Species Reactivity	Human
Specificity	Detects endogenous human Peroxiredoxin 1 in Western blots.
Source	Monoclonal Mouse IgG ₁ Clone # 477609
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human Peroxiredoxin 1 Met1-Lys199 Accession # Q06830
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Knockout Validated	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.
Immunoprecipitation	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Human Peroxiredoxin1 (Prx-1, also known as Thioredoxin peroxidase 2) is a 22 kDa antioxidant enzyme that belongs to the *typical* 2-Cys class of the THP/ahpC family of proteins. The molecule is 199 amino acids (aa) in length and has two catalytic cysteines, one at Cys52 and a second at Cys173. Prx-1 is an obligate homodimer. In its inactive state, Prx-1 is apparently noncovalently associated. Upon peroxide binding to Cys52 of subunit 1, the Cys173 of subunit 2 interacts with Cys52 of subunit 1 to complete the antioxidation, generating a disulfide bond between Cys52 and Cys173. Subsequent reduction restores the subunits to the basal state. There are apparently two additional isoforms; one shows a premature truncation after aa 171, while the second shows a deletion of aa 21-121. Human Prx-1 is 96% aa identical to mouse Prx-1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.