

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Periostin/OSF-2 in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 613618R
Purification	Protein A or G purified from cell culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Periostin/OSF-2 Met1-Gln836 Accession # Q15063
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

ELISA Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Periostin, also known as OSF-2, is a secreted matricellular protein with functions in extracellular matrix formation, cell migration, and inflammation (1). It is secreted as a 90 kDa monomer that can aggregate into >170 kDa higher-order multimers (2). Periostin contains an N-terminal EMI domain followed by four tandem FAS1 domains [takeshita]. Mature human Periostin shares 91% aa sequence identity with mouse and rat Periostin. Alternative splicing generates additional isoforms with various deletions in the C-terminal region following the FAS domains. Periostin is expressed by mesenchymal cells such as vascular smooth muscle cells, fibroblasts, osteoblasts, and odontoblasts in developing teeth (4-7). It is up-regulated in many carcinomas (2, 8). Periostin binds to Integrins $\alpha_v\beta_3$ and $\alpha_v\beta_5$ (2, 9), leading to enhanced cell adhesion and cell migration (2, 5, 6). It enhances Fibronectin and Collagen I production and promotes collagen fibrillogenesis (10, 11). It also induces epithelial-mesenchymal transition, tumor growth, invasion, and metastasis (9). Periostin induces the expression of VEGF R2 on endothelial cells and VEGF-C in tumor cells, and it can induce tumor lymphangiogenesis (8, 12). Periostin plays an important role in heart valve development and tissue healing after myocardial infarction (5, 13, 14). In asthma, it is up-regulated in bronchial epithelium and plays both destructive and protective roles by inducing eosinophil infiltration and inhibiting goblet cell metaplasia and mucus production, respectively (15, 16).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.