

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human β -1,4-Galactosyltransferase 1/B4GalT1 in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 394709
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human β -1,4-Galactosyltransferase 1/B4GalT1 Ala44-Ser398 Accession # P15291
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunocytochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

β 4GalT1 is one of seven β 1,4 galactosyltransferases that transfer galactose in a β 1,4 linkage to acceptor sugars including GlcNAc, and Glc, and Xyl. By sequence similarity, the β 4GalTs form four groups: β 4GalT1 and β 4GalT2, β 4GalT3 and β 4GalT4, β 4GalT5 and β 4GalT6, and β 4GalT7 (1). β 4GalT1 is unique among the seven enzymes because it can be expressed either as membrane associated form or secreted form (2). The secreted form is restricted to lactating mammary tissues where the enzyme forms a heterodimer with α -lactalbumin to catalyze the synthesis of lactose (3). The membrane form can reside either in the Golgi apparatus, where it adds galactose to N-acetylglucosamine residues, or on cell surface, where it functions as a recognition molecule during a variety of cell to cell and cell to matrix interactions, by binding to specific oligosaccharide ligands on opposing cells or in the extracellular matrix (4). The two enzymatic forms result from alternate transcription initiation sites and post-translational processing (5). Defects in β 4GalT1 are the cause of congenital disorder of glycosylation type 2D (CDG2D) (6). The amino acid sequence of human B4GALT1 is 88%, 87% and 71% identical to that of rat, mouse/canine and chicken.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.