

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse NTB-A/SLAMF6 in direct ELISAs and Western blots. In direct ELISAs, no cross-reactivity with recombinant human NTB-A or recombinant mouse CD84 is observed.
Source	Monoclonal Rat IgG _{2A} Clone # 448318
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse NTB-A/SLAMF6 Glu31-Asn239 Accession # Q9ET39
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

NTB-A, also known as Ly108 and SLAMF6, is a 60 kDa type I transmembrane glycoprotein that belongs to the SLAM subgroup of the CD2 family (1). Mature mouse NTB-A consists of a 209 amino acid (aa) ECD with one Ig-like V-type and one Ig-like C2-type domain, a 23 aa transmembrane segment, and an 89 aa cytoplasmic domain with two immunoreceptor tyrosine-based switch motifs ITSMs (2). Within the ECD, mouse NTB-A shares 48% and 70% aa sequence identity with human and rat NTB-A, respectively. The ECD of mouse NTB-A shares 20%-34% aa sequence identity with comparable regions of mouse 2B4, BLAME, CD2F-10, CD84, CD229, CRACC, and SLAM. An alternatively spliced isoform diverges after the second ITSM (2). NTB-A is expressed on the surface of NK, T, and B lymphocytes as well as eosinophils (3-5). It interacts homophilically through weak associations between the Ig-V type domains (5-7). NTB-A functions as an activating coreceptor on NK and T cells (3, 5, 6, 8). Tyrosine phosphorylation in the membrane proximal ITSM enables specific association with EAT-2, an interaction that is required for NTB-A mediated cytotoxicity of NK cells (9). Phosphorylation-dependent NTB-A association with SAP is required for full production of NK cell IFN-γ (5, 9). This interaction is independent of EAT-2 binding and appears to involve the membrane distal ITSM (5, 9). NTB-A deficient mice show weakened Th2 responses and elevated levels of neutrophil-derived inflammatory mediators (10). On B cells, NTB-A modulates immunoglobulin class switching and the balance between tolerance and autoimmunity (5, 11). The isoform with the divergent C-terminal tail is overexpressed in B cells from lupus-prone mice (11).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.