

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Brevican in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 450616
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Brevican isoform 1
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunocytochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Brevican, also called BEHAB, is a secreted member of the the lectican family of proteoglycans that share a common domain structure (1). Brevican contains an Ig-like V-set domain, two link domains, a Glu-rich region, a central region with glycosaminoglycan (GAG) modifications, an EGF-like domain, a C-type lectin domain, and a C-terminal Sushi/CRP-like domain (2). Brevican is restricted to the CNS and is expressed by astrocytes, oligodendrocytes, and neurons (3-7). A GPI-anchored alternate splice form exists that is truncated following the central (GAG) region (2, 8). Brevican is cleaved by multiple proteases and exists in a number of distinct fragments (5, 9, 10). Full-length brevican consists of a 97 kDa core protein with up to approximately 100 kDa of attached chondroitin sulfate but not heparan sulfate chains (4, 7, 11, 12). Brevican associates with the extracellular matrix, perineuronal nets, and astrocyte cell surfaces by means of its chondroitin sulfate, GPI anchor, hyaluronic acid-binding link domains, and the core protein (4, 7, 8, 13). The secreted isoform is dominant during brain development and is upregulated in astrocytes following brain injury (2, 14). In human and rat, an under-glycosylated form of brevican is upregulated in highly aggressive glioma but not in low-grade glioma or other brain pathologies (15, 16). In mouse and rat, levels of an ADAMTS4-generated 55 kDa N-terminal fragment increase during remodeling after excitotoxic injury (11, 12). Human brevican shares 90%, 80%, and 80% aa sequence identity with bovine, mouse, and rat brevican, respectively. Within the Ig-like and two link domains, brevican shares 45%-51% aa sequence identity with aggrecan, neurocan, and versican.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.