

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human IFN-α/β R2 in direct ELISAs.
Source	Monoclonal Mouse IgG _{2B} Clone # 493715
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line, NS0-derived recombinant human IFN-α/β R2 Ile27-Lys243 Accession # P48551
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

IFN-α/β R2, also known as IFNAR2, is a 100 kDa glycoprotein in the class II cytokine receptor family. These proteins form heterodimeric receptor complexes that transduce signals from the interferon, IL-10, and IL-28 families of cytokines (1, 2). IFN-α/β R2, in association with IFN-α/β R1, is required for mediating the antiviral, antiproliferative, and apoptotic effects of the type I interferons IFN-α and IFN-β. IFN-α/β R2 is the principal ligand binding subunit of the receptor. Ligand binding is stabilized by the subsequent association with IFN-α/β R1, resulting in the formation of a signaling ternary receptor complex (3, 4). Mature human IFN-α/β R2 consists of a 217 amino acid (aa) extracellular domain (ECD) with two fibronectin type III repeats, a 21 aa transmembrane segment, and a 251 aa cytoplasmic domain. Alternate splicing generates a secreted isoform that corresponds to the ECD and a 50 kDa transmembrane isoform with a substituted and truncated cytoplasmic region (5, 6). The short isoform is impaired in its ability to activate signaling molecules and functions as a dominant negative receptor subunit (7-9). IFN-α/β R2 is also subject to presenilin-dependent intramembrane proteolysis, resulting in the liberation of nearly the entire ECD as well as the cytoplasmic domain which migrates to the nucleus and can inhibit gene transcription (10). High concentrations of soluble IFN-α/β R2 bind and neutralize IFN-α and IFN-β, while lower concentrations prolong the antiviral activity of circulating IFN-β but not IFN-α (11). Human but not mouse IFN-α/β R2 constitutively associates with STAT4, which may account for species specific differences observed in type I interferon responses (12). Within the ECD, human IFN-α/β R2 shares 63%, 60%, and 48% aa sequence identity with bovine, mouse, and ovine IFN-α/β R2, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.