

## DESCRIPTION

<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human Integrin $\beta$ 4/CD104 in Western blots. In Western blots, no cross-reactivity with recombinant human Integrin $\beta$ 1, $\beta$ 2, $\beta$ 3, $\beta$ 5, $\beta$ 6, $\beta$ 8, or recombinant mouse Integrin $\beta$ 4 is observed.
<b>Source</b>	Monoclonal Mouse IgG <sub>2B</sub> Clone # 422325
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Chinese hamster ovary cell line CHO-derived recombinant human Integrin $\beta$ 4/CD104 Asn28-Ser710 Accession # P16144
<b>Conjugate</b>	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
<b>Formulation</b>	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Flow Cytometry</b>	0.25-1 $\mu$ g/10 <sup>6</sup> cells	A431 human epithelial carcinoma cell line

## PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Protect from light. Do not freeze.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul>

## BACKGROUND

Integrin  $\beta$ 4 (also GP150 and CD104) is a 150-200 kDa member of the Integrin beta family. It forms noncovalent heterodimers with Integrin  $\alpha$ 6 and participates in the formation of epithelial hemidesmosomes. Human Integrin  $\beta$ 4 is a type I transmembrane glycoprotein that is 1795 amino acids (aa) in length. It contains a 683 aa extracellular domain (ECD) (aa 28-710) and a large 1089 aa cytoplasmic region. Two ECD alternative splice forms may exist. One shows an 84 aa substitution for the 84 aa between aa 621-704. A second shows a 144 aa substitution for the first 704 amino acids. Human  $\beta$ 4 ECD shares 88% aa sequence identity with mouse  $\beta$ 4 ECD.

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.