

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse IL-7 in direct ELISAs.
Source	Monoclonal Rabbit IgG Clone # 2304B
Purification	Protein A or G purified from cell culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant mouse IL-7 Glu26-Ile154 Accession # P10168
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

IL-7, previously known as pre-B-cell growth factor and lymphopoietin-1, was originally purified on the basis of its ability to promote the proliferation of precursor B-cells. It has now been shown that IL-7 can also stimulate the proliferation of thymocytes, T cell progenitors and mature CD4⁺ and CD8⁺ T cells. IL-7 can induce the formation of lymphokine-activated killer (LAK) cells as well as the development of cytotoxic T lymphocytes (CTL). IL-7 was also shown to induce the V(D)J rearrangement of the T cell receptor β gene in mouse fetal thymocytes. Among myeloid lineage cells, IL-7 can up-regulate the production of pro-inflammatory cytokines and stimulate the tumoricidal activity of monocytes/macrophages. IL-7 is expressed by adherent stromal cells from various tissues.

Mouse IL-7 cDNA encodes a precursor protein of 154 amino residues containing a 25 amino acid residue signal peptide. Human IL-7 has approximately 65% amino acid sequence identity with mouse IL-7 and both proteins exhibit cross-species activity.

IL-7 bioactivities are mediated by the binding of IL-7 to functional high-affinity receptor complexes. The ligand binding subunit (IL-7 R) of the IL-7 receptor complex has been cloned from human and mouse sources. In addition to the membrane-anchored form of the IL-7 receptor, a human cDNA clone that encodes a soluble form of the IL-7 R has also been isolated. The γ chain of the IL-2 receptor complex has been shown to be an essential component for IL-7 signal transduction. Both IL-7 R and IL-2 R γ are members of the hematopoietin receptor superfamily. Cells known to express IL-7 receptors include pre-B cells, T cells and bone marrow cells.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.