

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human DMP-1 in ELISAs. In sandwich immunoassays, no cross-reactivity with recombinant mouse DMP-1, recombinant human (rh) CD44, and rhComplement Factor H is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 560716
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human DMP-1 isoform 2 Leu17-Tyr497 Accession # NP_001073380
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

ELISA Capture (Matched Antibody Pair)	Optimal dilution of this antibody should be experimentally determined.
ELISA Detection (Matched Antibody Pair)	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Dentin matrix protein 1 (DMP-1) is a member of the SIBLING family that also includes bone sialoprotein, dentin sialophosphoprotein, MEPE, and osteopontin. These highly phosphorylated integrin-binding proteins are rich in acidic amino acids and function in the formation of calcified bone and tooth matrix (1, 2). The phosphate content, spacing of acidic residues, and calcium-dependent dimerization of DMP-1 contribute to its ability to sequester calcium phosphate clusters and promote hydroxyapatite (HA) crystal formation (3-5). Rodent DMP-1 is cleaved by BMP-1 family proteases at a single site which is conserved in human, generating a 37 kDa N-terminal and a 57 kDa C-terminal fragment (6). The N-terminal fragment in rat carries chondroitin sulfate (7). The C-terminal fragment alone can nucleate HA crystals, while crystal growth into a needle-like morphology is inhibited by the N-terminal fragment (3, 4). Crystal maturation is dependent on the presence of type I collagen (4). DMP-1 is required for odontoblast differentiation as well as dentin formation (8). Nonphosphorylated DMP-1 is targeted to the nucleus, where it activates the transcription of odontoblast and osteoblast specific genes (9, 10). Early in osteoblast maturation, nuclear DMP-1 is extensively phosphorylated by casein kinase II, triggering its secretion (9). DMP-1 mutations in humans are associated with hypophosphatemia and FGF23 overexpression (11, 12). DMP-1 induces the activation of proMMP-9 and displaces mature MMP-9 from TIMP1 (13). DMP-1 tethering of MMP-9 to the cell surface via CD44 and integrins αvβ3 and αvβ5 promotes tumor cell invasiveness in vitro (14). Full length DMP-1 circulates in human serum in a tight complex with complement factor H (13, 14). When first bound to CD44 or integrin αvβ3, DMP-1 can anchor factor H to the cell surface and protect the cell from complement-mediated lysis (15). Mature human DMP-1 shares 61%-67% amino acid sequence identity with bovine, mouse, and rat DMP-1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.