

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human COX-2 in Western blots.
Source	Monoclonal Mouse IgG _{2B} Clone # 495222
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human COX-2 Ala18-Ser112 and Gln386-Leu604 Accession # P35354
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Cyclooxygenase-2 (COX-2) also known as prostaglandin G/H synthase 2 (PGHS2) is a 70 kDa microsomal enzyme that belongs to the prostaglandin G/H synthase family. It is inducibly-expressed by a number of cell types, including fibroblasts, vascular smooth muscle cells, endothelium, and monocytes. Functionally, COX-2 is a homodimer that catalyzes two steps in the conversion of arachadonic acid to prostaglandin H₂. Mature human COX-2 is 587 amino acids (aa) in length and contains one EGF-like domain (aa 18-55), a potential membrane interacting region (aa 277-292) and a globular catalytic domain (aa 293-604). At least one splice form exists that shows an 11 aa substitution for the C-terminal 451 amino acids. Over the amino acid range of the immunogen, human COX-2 shows 83% aa identity to mouse COX-2.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.