

| DESCRIPTION        |   |  |
|--------------------|---|--|
| Species Reactivity | Human   |  |
| Specificity        | Detects human CD20 in direct ELISAs.  |  |
| Source             | Recombinant Monoclonal Rabbit IgG Clone # 2709D   |  |
| Purification       | Protein A or G purified from cell culture supernatant   |  |
| Immunogen          | Human CD20 Peptide. The epitope recognized by rituximab<br>Binder M, et al. (2006) 108:1975-8. doi: 10.1182/blood-2006-04-014639. Epub 2006 May 16. PMID: 16705086                                      |  |
| Conjugate          | Alexa Fluor 647<br>Excitation Wavelength: 650 nm<br>Emission Wavelength: 668 nm   |  |
| Formulation        | Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide.  |  |
|                    | *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |  |

| APPLICATIONS                                    |   |
|---|---|
| Please Note: Optimal dilutions should be determ | ined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.              |
| Flow Cytometry                                  | Titration recommended for optimal concentration with starting range of 0.1-1 μg/1 million cells. Sample used for this experiment was Human PBMC |

| PREPARATION AND STORAGE |  |  |
|-------------------------|--|--|
| Shipping                | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.      |  |
| Stability & Storage     | <ul> <li>Protect from light. Do not freeze.</li> <li>12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul> |  |

## BACKGROUND

CD20 is a 297 amino acid (aa) non-glycosylated type III membrane protein that is the founding member of the MS4A tetraspanin protein family. It is expressed on pre-B, naïve and mature B lymphocytes and B-cell lymphomas. CD20 is associated with lipid rafts upon crosslinking, and appears to play a role in regulation of ion influx. The one extracellular segment of 48 as shows 65% as sequence identity between human and mouse CD20.

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

Rev. 12/1/2022 Page 1 of 1



**Global** bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL +1 612 379 2956 USA TEL 800 343 7475 Canada TEL 855 668 8722 China TEL +86 (21) 52380373 Europe | Middle East | Africa TEL +44 (0)1235 529449