

DESCRIPTION


Species Reactivity	Human
Specificity	Stains human CEACAM-8/CD66b transfectants but not irrelevant transfectants in flow cytometry.
Source	Recombinant Monoclonal Mouse IgG ₁ Clone # 913542
Purification	Protein A or G purified from cell culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human CEACAM-8/CD66b Gln35-His141 Accession # P31997
Conjugate	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the [Technical Information](#) section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

CEACAM-8 (Carcinoembryonic Antigen-related Cell Adhesion Molecule 8) also known as CD66b, CD67 and NCA-95, is a 90 kDa member of the CEACAM subfamily of the CEA family of proteins. It is expressed by neutrophils and eosinophils, and serves as a binding partner for CEACAM-6 and Galectin-3. Mature human CEACAM-8 is a 287 amino acid GPI-linked glycoprotein. It contains one V-type and two C2-type Ig-like domains. No definitive rodent CEACAM-8 has been reported.