

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects recombinant mouse (rm) CXCL17/VCC-1 in direct ELISAs and Western blots. In direct ELISAs, 50% cross-reactivity with recombinant human CXCL17 is observed/VCC-1.
Source	Monoclonal Rat IgG _{2B} Clone # 510614
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant mouse CXCL17/VCC-1 Ser31-Leu119 Accession # Q5UW37
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

CXCL17, also known as dendritic cell and monocyte chemokine-like protein (DMC) and VEGF-correlated chemokine-1 (VCC-1), is a secreted molecule with a size and predicted three-dimensional folding pattern similar to that of chemokines CXCL8/IL-8 and CXCL14/BRAK (1, 2). It has no predicted N-glycosylation sites, so cleavage of a 22 amino acid (aa) signal sequence likely results in a mature mouse CXCL17 molecule of 97 aa and 11 kDa size. CXCL17 is constitutively produced by airway and intestinal epithelium (1). It induces the chemotaxis of quiescent, but not LPS-activated peripheral blood monocytes and dendritic cells, and also binds these cells specifically (1). CXCL17 expression is increased in endothelial cells when they are induced to form tubes *in vitro* (2). Transgenic overexpression in NIH3T3 cells causes upregulation of proteins such as VEGF and FGF basic, and increases cell growth rate and tumorigenicity (2). CXCL17 and two other chemokines that play roles in angiogenesis, CXCL1/GRO and CXCL8/IL-8, show significantly correlated expression with that of VEGF in primary lung, breast and esophageal tumors (2). CXCL17 is therefore suggested to play a role in tumor angiogenesis. Mature mouse CXCL17 shares 82%, 71% and 66% amino acid sequence identity with rat, human and bovine CXCL17, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.