

Mouse CCL12/MCP-5 Alexa Fluor® 750-conjugated Antibody

Monoclonal Rat IgG_{2A} Clone # 63205 Catalog Number: FAB428S

100 µg

| DESCRIPTION | | |
|--------------------|---|--|
| Species Reactivity | Mouse | |
| Specificity | Detects mouse CCL12/MCP-5 in ELISAs and Western blots. | |
| Source | Monoclonal Rat IgG _{2A} Clone # 63205 | |
| Purification | Protein A or G purified from hybridoma culture supernatant | |
| Immunogen | E. coli-derived recombinant mouse CCL12/MCP-5 Gly23-Gly104 Accession # Q62401.1 | |
| Conjugate | Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm | |
| Formulation | Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide | |
| | *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. | |

| APPLICATIONS | | | |
|---|--|--|--|
| Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website. | | | |
| ELISA Capture (Matched Antibody Pair) | Optimal dilution of this antibody should be experimentally determined. | | |
| ELISA Detection (Matched Antibody Pair) | Optimal dilution of this antibody should be experimentally determined. | | |
| Neutralization | Optimal dilution of this antibody should be experimentally determined. | | |
| Western Blot | Optimal dilution of this antibody should be experimentally determined. | | |

| PREPARATION AND STORAGE | | |
|-------------------------|---|--|
| Shipping | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. | |
| Stability & Storage | Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied | |

BACKGROUND

CCL12 is a cloned mouse CC chemokine most closely related to human MCP-1 (66% amino acid (aa) sequence identity in the mature protein). Mouse CCL12 encodes a 104 aa residue precursor protein with a 22 aa residue predicted hydrophobic signal sequence that is cleaved to generate a 82 aa residue mature protein. CCL12 is expressed constitutively in the thymus and lymph nodes. Under inflammatory conditions, CCL12 expression is also induced in activated macrophages and mast cells. The gene for mouse MCP-1 has been mapped to the CC chemokine cluster on chromosome 11. Recombinant CCL12 has been shown to be a potent chemoattractant for monocytes and lymphocytes but not neutrophils. At high concentrations, CCL12 will also chemoattract eosinophils. CCL12 has been found to be a functional ligand for CCR2 but not CCR1, CCR3, or CCR5.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

Rev. 9/21/2025 Page 1 of 1

China | info.cn@bio-techne.com TEL: 400.821.3475